



Contents lists available at ScienceDirect

Reproductive Toxicology

journal homepage: www.elsevier.com/locate/reprotox



Evaluating chemical effects on mammary gland development: A critical need in disease prevention

Gwendolyn Osborne^a, Ruthann Rudel^b, Megan Schwarzman^{a,*}

^a Center for Occupational and Environmental Health, University of California at Berkeley, Berkeley, CA, USA

^b Silent Spring Institute, Newton, MA, USA

ARTICLE INFO

Article history:

Received 26 November 2013
Received in revised form 26 June 2014
Accepted 22 July 2014
Available online xxx

Keywords:

Mammary gland
Development
Breast cancer
Endocrine disrupting compounds
Toxicity testing

ABSTRACT

Although understanding the environmental factors that contribute to breast cancer could improve disease prevention, standard chemical testing protocols do not adequately evaluate chemicals' effects on breast development. Evidence suggests: (1) mammary gland (MG) development is a complex process that extends from gestation through fetal and neonatal growth, puberty, and pregnancy; (2) altered MG development can increase the risk of breast cancer and other adverse outcomes; and (3) chemical exposures during susceptible windows of development may alter the MG in ways that increase risk for later disease. Together, these highlight the need to better understand the complex relationship between exposure to endocrine disrupting compounds (EDCs) and the alterations in MG morphology and gene expression that ultimately increase disease risk. Changing guideline toxicity testing studies to incorporate perinatal exposures and MG whole mounts would generate critical knowledge about the effects of EDCs on the MG and could ultimately inform disease prevention.

© 2014 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/3.0/>).

1. Introduction

Breast cancer is the most common cancer in women worldwide and the second leading cause of cancer death in American women, after lung cancer [1,2]. The probability of a woman being diagnosed with breast cancer during her lifetime is one in eight [3]. While inherited risk factors explain up to a third of breast cancer cases [4,5], the risk factors for the remaining majority of cases are not well understood. In addition to a steady increase in female breast cancer, the incidence of male breast cancer has increased in the past few decades in the U.S. and internationally [6]. The correlation of male and female incidence rates over time suggests that there may be risk factors that are similar for both men and women [6,7]. Exposure to chemicals in the environment and consumer products is hypothesized to contribute to cancer risk [8–10], as well as to other breast health outcomes, such as impaired lactation [11] or male gynecomastia [12].

Despite emerging evidence that chemical exposure may contribute to breast cancer risk, most chemicals are not evaluated for their potential impact on breast tissue, particularly during vulnerable stages of development. Although understanding the environmental factors that contribute to breast cancer has

the potential to dramatically improve prevention of the disease, standard chemical testing protocols do not adequately evaluate chemicals' effects on breast tissue. Most protocols lack an evaluation of the effects of exposures during critical periods of mammary gland (MG) development, as well as an assessment of functional outcomes such as lactation impairment [13].

As a result, we have only limited evidence of how chemicals such as endocrine disruptors (EDs) alter MG development, and how those effects contribute to adverse outcomes later in life. The evidence that exists suggests three key points: (1) MG development is a complex process that extends from gestation through multiple life stages; (2) altered MG development can increase the risk of breast cancer, as well as other adverse outcomes; and (3) chemical exposures during susceptible windows of development may alter the MG in ways that increase risk for later disease. Together, these points highlight the need to better understand the complex relationship between environmental exposures and the alterations in MG morphology and gene expression that ultimately increase the risk of disease.

2. Mammary gland development is a complex process extending from gestation through multiple life stages

Normal breast development in both humans and rodents consists of a series of well-orchestrated events that are finely regulated by a balance of hormones, growth factors, and

* Corresponding author. Tel.: +1 5106434685; fax: +1 510 642 5815.
E-mail address: mschwarzman@berkeley.edu (M. Schwarzman).

stromal factors [14,15]. Growth depends on endocrine signaling from the hypothalamic–pituitary–gonadal axis, autocrine and paracrine hormones, and growth factors from outside tissues. Crosstalk between the epithelium and surrounding stroma also helps to balance proliferation and apoptosis during normal developmental remodeling of the MG [15,16].

The MG is distinct from other tissues in that it undergoes a significant portion of its development postnatally; in addition to the fetal/neonatal period, puberty and pregnancy are critical periods of MG development. In most mammals, MG development begins with the formation of the mammary, or milk, line. This separates into individual placodes, each of which develops into a ductal tree that embeds in a fat pad to form the mammary bud [15,17,18]. Factors that interfere with signaling from the surrounding fat pad to the mammary bud can potentially alter the timing of development or formation of glandular structures [19,20].

Subsequent to the neonatal period there is little epithelial growth until puberty. During puberty, mammary growth is exponential; this is a period of several weeks in rodents, or years in girls, during which the fat pad rapidly fills with epithelial cells to become the adult form of the gland. The epithelium develops bundles of ducts, which then form club-like structures, called terminal end buds (TEB) in humans. Each TEB cleaves into alveolar buds and sprouts into ductules. This structural unit, comprised of the terminal duct and the ductules, is called the terminal ductal lobular units (TDLU) [21]. Rodents also form TEBs, which are the structures most functionally equivalent to TDLUs in humans. These teardrop-shaped structures are the sites of future ductal branching and disappear as the gland differentiates [20].

In both humans and rodents, the key periods of development in MG maturation are regulated by the activation (in the fetal and neonatal periods) and later the reactivation (during puberty) of the hypothalamic–pituitary–gonadal (HPG) and hypothalamic–pituitary–adrenal (HPA) axes, which control the release of hormones [14,16,18].

The gland reaches a fairly static state from first menstruation until a pregnancy occurs. During pregnancy, the gland undergoes another period of rapid differentiation, involving branching and the development of lobulo-alveoli to prepare for lactation [19].

Male breast development also occurs in utero, but the androgen surge halts further development immediately prior to birth. Exposure to anti-androgens has been shown to lead to retained nipples in male rats [22–24].

3. Alterations to mammary gland development can increase the risk of breast cancer and other adverse outcomes

3.1. Human evidence

The fetal origins of adult disease hypothesis propose that disturbances to the fetal environment have physiological and structural consequences with the potential to alter an individual's disease risk as an adult [25]. Human epidemiological studies provide support for this hypothesis with respect to breast health outcomes, as alterations to the finely regulated process of MG development have been shown to produce changes that affect women later in life. For example, various birth parameters have been associated with breast cancer risk. Birth weight [26], longer birth length [27], older maternal and paternal age [28], and in utero exposure to synthetic estrogen and diethylstilbestrol (DES) [29] have all been associated with an increased risk of later breast cancer, while maternal preeclampsia is associated with a lower risk [28,30]. Each of these factors affects the earliest period of breast development that starts before birth.

The second period of breast development occurs during puberty. Ionizing radiation has the greatest effect on later breast cancer risk when exposure occurs before the age of 20 [31,32], suggesting that the period from childhood through adolescence is another significant period of vulnerability for the human breast.

Pregnancy is a third period of breast development during which external factors can alter disease susceptibility. Younger age at first birth, multiple gestation pregnancies (twins and greater), preeclampsia, pre-pregnancy obesity, and gestational hypertension may all lower maternal breast cancer risk (reviewed by [33,34]), while increased placental weight [35], higher infant birth weight [34], and DES exposure during pregnancy are associated with increased risk of maternal breast cancer. One factor driving these changes is total lifetime exposure to ovarian hormones. Lower cumulative exposure to estrogen – such as with pre-eclampsia – seems to protect against breast cancer [28]. Higher exposure to progesterone may increase risk of breast cancer, and affecting progesterone or progesterone receptor signaling pathways promotes breast cancer progression [36,37].

The relationship between lifetime estrogen and progesterone exposure and breast cancer risk provides a framework for understanding how chemicals that affect hormone homeostasis may alter breast development and ultimately cancer risk.

3.2. Rodent evidence

The rodent MG undergoes staged development analogous to that observed in humans during gestation, puberty, and pregnancy. Animal evidence further supports the hypothesis that there are periods of vulnerability during breast development that influence later life outcomes. A structure particularly important in rodent MG development and carcinogen susceptibility is the TEB [20,38]. The TEB has the greatest number of proliferating cells and the shortest cell cycle of the structures in the developing MG. Malignant tumors, such as adenocarcinomas induced by certain carcinogens, are most common in rodents following exposures that occur between days 40 and 46 of life (correlating with puberty in humans), the period of development when TEBs are most actively differentiating into alveolar buds. Benign tumors, such as adenomas, fibroadenomas, and mammary cysts, are thought to arise from the more differentiated alveolar buds [39].

Anything that changes the timing of mammary development will affect the timing of the presence of TEBs, and therefore the window of susceptibility to carcinogens. Earlier induction of MG development in rodents leads to a greater number of TEBs compared to terminal ducts and increased alveolar budding at the time of weaning, followed by the development of more lobules than in control animals [40,41]. On the other hand, late initiation of mammary development causes decreased longitudinal growth of the epithelium and fewer TEBs, and decreased alveolar budding at weaning [42]. As development progresses, these glands may have more TEBs at puberty, because the pace of development is slower [43]. It is hypothesized that factors that lengthen the period when TEBs are present lengthen the period during which the MG is susceptible to carcinogens.

3.3. Non-cancer effects

Altered MG development is also associated with non-cancer effects such as lactation impairment and gynecomastia in both rodents and humans. While it is recommended that infants are breastfed exclusively for at least the first six months of life [44], several million mothers are unable to breastfeed or have significant difficulty breastfeeding each year [45]. Research in rodents suggests that factors that interfere with MG growth and differentiation can negatively affect both the gland's ability to produce milk

and the milk composition [45,46]. Factors that accelerate MG development also make the gland less sensitive to ovarian and pituitary hormones. Because effective lactation depends on these hormonal signals to stimulate the gland, a less responsive gland is less likely to be capable of normal milk production [43].

Gynecomastia, or benign breast enlargement due to proliferation of the ductal epithelium [12], occurs in up to two-thirds of pubertal boys [47]. It is hypothesized that the altered MG development that causes this condition arises due to an imbalance in estrogen and androgen [48,49].

4. Exposure to chemicals during windows of susceptibility can alter mammary gland structure and function, increasing cancer risk

4.1. The role of hormones and EDs

Many of the established risk factors for breast cancer are proxy measures for changes in the hormonal environment [50–52]. For example, high birth weight, birth length, and placental weight are associated with increased endogenous intrauterine estrogen exposure [53,54], while pre-eclampsia is associated with increased androgen levels [55]. Furthermore, DES, a synthetic estrogen prescribed to prevent miscarriage from the 1940s through 1970s, has been associated with breast cancer in women who were directly exposed, as well as their daughters [29,56].

Levels of circulating estrogens and androgens are associated with breast cancer risk in both premenopausal and postmenopausal women [57–59]. Breast cancer risk is elevated by extended exposure to high levels of endogenous hormones, which can occur with obesity [60] or as a result of early age at menarche or late age of first pregnancy and menopause [61,62]. However, the relationship between hormones and breast cancer is complicated by the fact that breast cancer is a heterogeneous disease with at least five molecular subtypes. The presence or absence of biological markers, such as estrogen receptors (ER+/ER–), progesterone receptors (PR+/PR–), and human epidermal growth factor receptor 2 (HER2+/HER2–), serve as approximations of the molecular subtypes [63]. The majority of information pertaining to hormonal risk factors is specifically associated with the ER+/luminal A subtype. The influence of hormones on ER– and triple negative (ER–, PR–, and HER2–) breast cancers are not as well characterized [57,64].

The endocrine system governs hormone production, and as such, plays a critical role in development and ongoing function of multiple organ systems. Disruption of the endocrine system is linked to multiple adverse outcomes, including metabolic disturbances (e.g., obesity) [65], and alterations to lactation, breast density [66], immune function [67,68], and the timing of puberty [69] and menopause [70]. An ED was originally defined as “an exogenous agent that interferes with the production, release, transport, metabolism, binding, action or elimination of natural hormones in the body responsible for the maintenance of homeostasis and the regulation of developmental processes” [71]. This definition has been simplified by Zoeller et al. [72] to define an ED as “an exogenous chemical, or mixture of chemicals, that interferes with any aspect of hormone action”. Compounds that have estrogen-like activity have been studied the most, yet chemicals with anti-estrogenic, androgenic, anti-androgenic, progesterone-like, and thyroid-like activity are also EDs [73].

4.2. EDs as carcinogens

Developmental exposure to EDs may contribute indirectly to a variety of adverse outcomes by altering MG development in ways that raise the risk of later disease. For example, chemical

alterations to breast development in early life can increase the tissue's susceptibility to a subsequent carcinogenic exposure, a so-called ‘second hit’ [43]. Carcinogen challenge models in experimental animals have demonstrated these effects by using a known chemical carcinogen following an early life chemical exposure to show the increased rate of tumor formation in response to the second exposure, relative to tumor incidence from the known carcinogen alone [10]. This effect has been observed in rodents with prenatal exposure to both estrogens and androgens [74–76]. Additionally, increased tumor number and accelerated tumor development after carcinogen challenge have been observed following early life exposure to known EDs such as alcohol [77], dioxin [78–80], bisphenol A (BPA) [81], and the phytoestrogen genistein [82,83].

Research suggests that, in addition to increasing susceptibility to a second hit by a chemical carcinogen, EDs may have direct carcinogenic effects. For example, DES causes cervical, vaginal, uterine, endometrial, and breast cancer via a combination of genotoxic, estrogen-receptor-mediated, and epigenetic effects, independent of subsequent carcinogen exposure [84]. A recent study demonstrated that BPA caused a non-significant increase in atypical hyperplasia and adenocarcinoma of the MG at doses ranging from 0.25 to 250 micrograms/kg-day without later exposure to another chemical carcinogen [85]. Taken together, these findings suggest that in some cases EDs may be complete carcinogens, not requiring a second hit to cause cancer.

4.3. Rodent evidence

Rodent studies suggest that many mammary carcinogens also alter MG development, often by accelerating or delaying development and/or alter branching and epithelial structure (specific chemicals reviewed in [10]). The effects of ED exposure are highly dependent on the level and timing of exposure, and can even have protective effects when exposure occurs at the right level and at a particular developmental stage. EDs can stimulate growth, causing the MG to mature with an increased ratio of fully differentiated structures compared to immature or undifferentiated structures. They can also reduce the ratio of proliferation to apoptosis in the epithelium [43], reducing the risk for tumor formation. For example, prenatal [78] and neonatal [79] dioxin exposure alters rodent MG differentiation, increasing MG tumor development following treatment with dimethylbenz[*a*]anthracene (DMBA). In contrast, dioxin exposure in long-term feeding studies decreases spontaneous tumor development [86], and dioxin exposure during pregnancy followed by treatment with DMBA four weeks later delays tumor formation compared to controls [87]. Likewise, genistein exposure during the peripubertal period decreases tumors after a carcinogen challenge, while perinatal and lifelong exposure appears to increase tumors in rats [88].

4.4. Human evidence

As with rodent studies, human studies demonstrate that outcomes depend on the timing of exposure. Cohn et al. [89] demonstrated that breast cancer is associated with DDT exposure, but only when exposure occurred before 14 years of age. This study used blood samples obtained before DDT use was banned to find that women who had high levels of serum DDT and were under 14 years old at time of exposure had a 5-fold increased risk of breast cancer, while women who were not exposed prior to 14 years of age demonstrated no association. Because puberty is a period of rapid development when the MG is particularly vulnerable to the effects of chemical exposure, lengthening puberty extends this window of susceptibility, thus increasing the possibility of ED exposure and adverse outcomes. This is particularly concerning in light of the

current shift toward earlier onset breast development in American and European girls, with girls now beginning breast development (thelarche) earlier and taking longer to reach full breast development than in previous generations [90]. Endogenous hormones play a central role in the onset and regulation of puberty; estrogen, the primary hormone in early breast development, promotes growth of the ducts, while progesterone promotes lobuloalveolar development [73,91]. The physical changes seen in premature thelarche may reflect the onset of pubertal maturation, or they may reflect exposure to endogenous or exogenous hormones independent of the maturity stage of reproductive or adrenal systems (pseudo-puberty) [92]. In general, precocious puberty is associated with various adverse outcomes, including metabolic syndrome and polycystic ovarian syndrome, behavioral disorders, testicular and prostate cancer, and breast cancer in later life [92]. While it is unknown if early thelarche is specifically associated with later breast disease, prolonging this sensitive developmental period increases the potential that susceptible tissues will be exposed to EDs or carcinogens [43].

4.5. The question of dose

In addition to timing, effects of ED exposure depend on dose: EDs can have complex dose response relationships – including showing opposite effects at lower doses relative to higher doses – which traditional toxicity testing methods are unable to detect [93]. Traditional toxicological studies typically use high doses to determine the maximum tolerated dose, lowest observed adverse effect level, and no observed adverse effect level, and extrapolate to estimate effects at lower doses. Developing tissues in particular can be exquisitely sensitive to very low doses of EDs. BPA, atrazine [94], PFOA [95], and dioxin [80] have each been shown to affect the MG at doses below those typically used in toxicological studies conducted for regulatory purposes. Exposure to concentrations of BPA expected to produce circulating free BPA levels above nanomolar concentrations has been shown to alter MG development, gene and protein expression, histogenesis, or to induce mammary hyperplasia in over a dozen studies (reviewed in [93]). Mammary epithelial cells exposed to nanomolar concentrations of BPA in culture demonstrate consistent effects [96].

Nonmonotonic dose-response (NMDR) curves are characterized by a nonlinear relationship between dose and effect, with the slope of the curve changing sign at some point in the range of doses examined. The effects of 17β -estradiol [97,98] and DES [99] on the MG have displayed NMDR curves in mice. Observations of NMDR curves in animal studies are particularly concerning because they raise questions about whether effects observed at higher doses can be extrapolated to lower doses more typical of human exposures, and raise concerns about missing important effects if testing is only performed at high doses.

4.6. Non-cancer effects

Finally, EDs that alter MG development may also affect the structure or function of the gland, causing changes such as gynecomastia in men or impaired lactation in women. In rodents, high doses of atrazine [100,101] and PFOA [95,102] have been shown to severely inhibit mammary development, which then affects lactation and impairs the growth of developing offspring. While these high doses are rarely seen in humans, it is possible that ecological systems may be exposed to levels that alter animals' development, impair lactation, and reduce offspring survival [43]. Other animal studies demonstrate that gestational exposure to dioxin [45,103], BPA [46,104–107], genistein [108], and Ziracin [109] not only alter mammary epithelial development, but impair lactation. It is possible that exposure to EDs contributes to

difficulty breastfeeding in women, but this outcome has not been investigated.

A limited number of epidemiological studies of prepubertal gynecomastia suggest an association with exposure to estrogenic or androgenic substances in boys [110,111]. Case reports have linked prepubertal gynecomastia to uses of tea tree oil and lavender oil, both of which have weak estrogenic and antiandrogenic activity [111]. While the mechanism is unclear, a variety of factors have been associated with gynecomastia in humans and domestic animals, including estrogen, androgen deficiency, and exposure to pyrethroid pesticides and some pharmaceuticals [48,112]. Since the growth of MG ductal epithelium is stimulated by estrogens and inhibited by androgens, EDs that alter the balance of estrogen or androgen stimulation may contribute to this condition [48]. Additionally, gynecomastia is frequently associated with other conditions that are related to hormonal imbalance. Vandenberg et al. [12] demonstrated that MG development was altered in male mice exposed to BPA during the perinatal period and studies of ethinyl estradiol and genistein show that the male MG is a highly sensitive marker of endocrine disruption [113], thus it appears that the male breast is also sensitive to EDs during susceptible periods of development.

5. Knowledge gaps

5.1. Understanding breast development

The three periods (i.e., fetal and neonatal growth, puberty, and pregnancy) of rodent MG development are known to be analogous to the periods of human breast development, however, research is needed to further describe the timing and characteristics of breast epithelial development and their relationship to adverse outcomes such as increased susceptibility to carcinogens or diminished lactation in both humans and rodents. Additionally, there are large gaps in our understanding of how MG development is regulated. It is important to understand what specific regulatory factors and genes are involved, as well as which receptor populations are present, and when, in order to understand the mechanisms through which these processes can be disrupted [55,73]. Elucidating the mechanisms involved in breast development would improve our understanding of breast cancer etiology, as well as non-cancer effects, such as lactation impairment and gynecomastia.

5.2. Understanding implications of altered development

Furthermore, while clear associations link various birth parameters to later life disease, the specific mechanisms through which perturbations to breast development cause disease or dysfunction years later are unknown. Understanding these mechanisms requires elucidation of the steps in mammary development and the processes that are susceptible to alteration. For example, epidemiologic evidence associates breast density with breast cancer risk. Density can be influenced by factors that are also associated with breast cancer risk (e.g., parity, number of births, and menopausal status) [114]. The stromal matrix of the breast tissue actively participates in the control of tumor growth [115–117] and may be a crucial target for carcinogens [118], suggesting that altered tissue architecture may be at the core of carcinogenesis. Research is needed to better understand the biological mechanisms responsible for these links and how perturbing them influences risk of cancer.

5.3. Understanding EDs' role in altered development and disease

Finally, animal studies demonstrate associations between ED exposure, altered MG development, and adverse outcomes.

However, the mechanisms by which EDs alter development and produce tumors are generally poorly understood. More information is needed on the types of chemicals, the doses, and the timing of exposure, as well as the ways that EDs act together, and the complex interactions among genetics, stress, diet, and endogenous and exogenous hormones.

Not only are there gaps in the knowledge concerning chemical causes of breast cancer, but the effects of ED on breast milk production and gynecomastia are largely unknown. Lactation impairment is usually identified by decreased pup weight or survival in multi-generational rodent studies. However, these surrogate markers do not adequately detect functional outcomes because they are not sensitive or specific for impaired lactation [13]. Additionally, multi-generational studies that can detect these effects are expensive and rare. An animal model has yet to be established to study gynecomastia, and there are very few studies investigating this condition.

6. Recommendations: improving test protocols to investigate how chemicals increase breast cancer risk and alter mammary gland structure and function

6.1. Include early developmental exposure in standard chemical testing protocols

While many knowledge gaps persist, recent research supports theories of development and later life outcomes in which environmental factors significantly alter phenotype [52]. Human epidemiologic studies are often unable to demonstrate associations between EDs and breast disease because exposures are not typically evaluated during critical periods of development. Longitudinal cohorts are of limited use in determining which early life exposures or factors increase breast cancer risk due to the difficulties of accurately measuring exposures and the length of time needed to see effects in humans. For example, it took 60 years for the effects of DES to appear [29]. As a result, the effects of EDs on the MG cannot be truly understood without animal studies. Such studies can be designed so that each animal model provides information concerning a particular aspect of human breast cancer and is able to represent a specific subset of the varieties of human breast disease [119].

In order to fully understand breast development and human breast disease, a variety of laboratory animal models are needed. Gaps in assessment of MG effects in the rodent toxicology guidelines produced by U.S. EPA, OECD, and NTP indicate that a paradigm shift is needed for these guideline studies to be able to detect critical alterations to MG development [13]. While the two-year carcinogenicity bioassays can assess MG histopathology, not all guidelines require exposure to the developing rodent. Developmental toxicity studies, which could provide information on developmental effects of EDs on the MG, typically do not evaluate MG tissue. Similarly, the standard one- or two-generation reproduction study could provide useful information on MG developmental effects but is not routinely used for this purpose.

The value of using the rat MG to elucidate potential modes of action of known mammary carcinogens has been shown by several investigators [119]. In most cases, these studies have been conducted in specialized study designs with treatment periods that replicate the developmental processes known to be important in the development of cancer. It is important to note that these inquiries and discoveries have not been conducted within the context of the standard 2-year carcinogenicity bioassay, as the usual protocols do not begin dosing until early adulthood. Reasons for this include (a) 2-year studies do not adequately assess changes that occur during especially sensitive life stages; (b) the studies are not designed to assess the consequences of early exposures that may

change tissue susceptibility to later exposures; and (c) the tests do not assess low-dose exposures to mixtures of chemicals that better approximate actual human exposure to chemicals found in the environment [120].

6.2. Evaluate MG development using whole mounts

Changing the two-year carcinogenesis study, developmental toxicity study, and multigenerational toxicity study guidelines would address these problems and improve our ability to understand ED effects on the MG. A protocol for evaluating whole mounts of MGs needs to be developed and validated that includes offspring exposed during perinatal and pubertal stages and that ensures outcomes are consistently reported among different laboratories. A whole mount is a tissue preparation technique that enables evaluation of the entire unsectioned organ [121]. This technique can identify morphological changes, as well as the temporal and spatial progression of epithelial development because it enables visualization of ductal branching [43]. Many laboratories use tissue sections, which alone are insufficient to detect developmental changes. Ideally, data from histological analyses could be used in combination with early life developmental end points seen in whole mounts to assess the later life impacts of early developmental changes [122]. Data generated from whole mounts could inform further investigations into the lactational impairment in offspring or tumor susceptibility using carcinogen-challenge protocols.

6.3. Other tools to improve testing

Another potential tool for evaluating chemical effects on the MG is the use of the male rat, which has demonstrated more sensitivity than other ED endpoints in male or females. A range of studies [113,123–125] indicate that the male rat MG is very sensitive to estrogen and may serve as a model for investigating the effects of EDs, although relevance for the female MG would have to be established [113,126].

Additional information can be gleaned from early biomarkers associated with MG cancer that can be measured using MG tissue blocks, serum samples, and other stored tissues. These include changes in levels of hormones or hormone receptors, receptor sensitivity, hyperplasia, gene markers, immunohistochemical markers, MG-specific gene markers, and effects on stromal–epithelial interactions [10]. Understanding these mechanisms would improve our knowledge of the relationship between altered MG development and later life outcomes.

Finally, toxicity information needs to be applicable to human-relevant endpoints. This can be achieved by including more time points during the investigation of MG developmental stages along with blood collection for dosimetry to understand the transition from normal mammary tissue to tumor development [43,73]. Careful characterization of internal dose is needed in all (not just guideline) studies [10,73]. Unlike laboratory animals, humans are exposed to many different environmental chemicals simultaneously, thus an understanding of biologically relevant mixtures that affect the mammary tissue is an important area of research.

7. Conclusion

Experimental evidence in rodent models and human epidemiological studies provide preliminary evidence that some EDs can affect normal MG development and function, and so may increase breast cancer risk. These clues indicate that traditional toxicology tests are likely missing many important effects on the MG. Chemical testing protocols should be standardized to require dosing during critical periods of development, assessment of MG development, structure, and function, and an assessment of the male MG.

A greater understanding of these mechanisms will help to clarify the risks of environmental exposures, provide evidence to reduce exposure, and ultimately reduce the burden of disease.

Conflict of interest statement

All authors confirm that they have no actual or potential competing interests regarding the submitted article.

Transparency document

The [Transparency document](#) associated with this article can be found in the online version.

Acknowledgements

This commentary grew out of work initiated in the Breast Cancer and Chemicals Policy Project, funded by the California Breast Cancer Research Program, grant number 15QB-8001. For the list of participants in that project, see <http://coeh.berkeley.edu/greenchemistry/cbcrp/htm>. The authors wish to thank Dr. Sarah Janssen for her review of the manuscript.

References

- [1] CDC. Invasive Cancer Incidence – United States, 2009. Atlanta, GA: Centers for Disease Control and Prevention; 2013.
- [2] Forouzanfar MH, Foreman KJ, Delossantos AM, Lozano R, Lopez AD, Murray CJL, et al. Breast and cervical cancer in 187 countries between 1980 and 2010: a systemic analysis. *Lancet* 2011;378(9801):1461–84.
- [3] Siegel R, Naishadham D, Jemal A. Cancer statistics, 2013. *CA Cancer J Clin* 2013;63(1):11–30.
- [4] American Cancer Society. Breast Cancer Facts & Figures 2011–2012. Atlanta, GA: American Cancer Society, Inc.; 2012.
- [5] Lichtenstein P, Holm NV, Verkasalo PK, Iliadou A, Kaprio J, Koskenvuo M, et al. Environmental and heritable factors in the causation of cancer—analyses of cohorts of twins from Sweden, Denmark, and Finland. *N Engl J Med* 2000;343(2):78–85.
- [6] Ly D, Forman D, Ferlay J, Brinton LA, Cook MB. An international comparison of male and female breast cancer incidence rates. *Int J Cancer* 2013;132(8):1918–26.
- [7] Anderson WF, Jatoi I, Tse J, Rosenberg PS. Male breast cancer: a population-based comparison with female breast cancer. *J Clin Oncol* 2010;28(2):232–9.
- [8] Brody JG. Everyday exposures and breast cancer. *Rev Environ Health* 2010;25(1):1–7.
- [9] Rudel RA, Attfield KR, Schifano JN, Brody JG. Chemicals causing mammary gland tumors in animals signal new directions for epidemiology, chemicals testing, and risk assessment for breast cancer prevention. *Cancer* 2007;109(12):2635–66.
- [10] Rudel RA, Fenton SE, Ackerman JM, Euling SY, Makris SL. Environmental exposures and mammary gland development: state of the science, public health implications, and research recommendations. *Environ Health Perspect* 2011;119(8):1053–61.
- [11] Neville MC, Walsh CT. Effects of xenobiotics on milk secretion and composition. *Am J Clin Nutr* 1995;61(3 Suppl.):687S–94S.
- [12] Vandenberg LN, Schaeberle CM, Rubin BS, Sonnenschein C, Soto AM. The male mammary gland: a target for the xenoestrogen bisphenol A. *Reprod Toxicol* 2013;37:15–23.
- [13] Makris SL. Current assessment of the effects of environmental chemicals on the mammary gland in guideline rodent studies by the U.S. Environmental Protection Agency (U.S. EPA), Organisation for Economic Co-operation and Development (OECD), and National Toxicology Program (NTP). *Environ Health Perspect* 2011;119(8):1047–52.
- [14] Hennighausen L. Molecular mechanisms of hormone controlled gene expression in the breast. *Mol Biol Rep* 1997;24:169–74.
- [15] Robinson GW. Cooperation of signaling pathways in embryonic mammary gland development. *Nat Rev Genet* 2007;8(12):963–72.
- [16] Brisken C, O'Malley B. Hormone action in the mammary gland. *Cold Spring Harb Perspect Biol* 2010;2:a003178.
- [17] Hennighausen L, Robinson GW. Think globally, act locally: the making of a mouse mammary gland. *Genes Dev* 1998;12:449–55.
- [18] Russo J, Russo IH. Development of the human breast. *Maturitas* 2004;49(1):2–15.
- [19] Hennighausen L, Robinson GW. Information networks in the mammary gland. *Nat Rev Mol Cell Biol* 2005;6(9):715–25.
- [20] Fenton SE. Endocrine-disrupting compounds and mammary gland development: early exposure and later life consequences. *Endocrinology* 2006;147(6 Suppl.):S18–24.
- [21] Russo J, Russo IH. The breast as a developing organ. *Mol Basis Breast Cancer* 2004;11–48.
- [22] Buckley J, Willingham E, Agras K, Baskin LS. Embryonic exposure to the fungicide vinclozolin causes virilization of females and alteration of progesterone receptor expression in vivo: an experimental study in mice. *Environ Health* 2006;5(4):1–6.
- [23] Gray LE, Ostby J, Furr J, Wolf CJ, Lambright C, Parks L, et al. Effects of environmental antiandrogens on reproductive development in experimental animals. *Hum Reprod Update* 2001;7(3):248–64.
- [24] Monosson E, Kelce WR, Lambright C, Ostby J, Gray Jr LE. Peripubertal exposure to the antiandrogenic fungicide, vinclozolin, delays puberty, inhibits the development of androgen-dependent tissues, and alters androgen receptor function in the male rat. *Toxicol Ind Health* 1999;15(1–2):65–79.
- [25] Barker DJP, Gluckman PD, Godfrey KM, Harding JE, Owens JA, Robinson JS. Fetal nutrition and cardiovascular disease in adult. *Lancet* 1993;341:938–41.
- [26] Michels KB, Xue F. Role of birthweight in the etiology of breast cancer. *Int J Cancer* 2006;119(9):2007–25.
- [27] Silva Idos S, De Stavola B, McCormack V, Collaborative Group on Pre-Natal Risk Factors and Subsequent Risk of Breast Cancer. Birth size and breast cancer risk: re-analysis of individual participant data from 32 studies. *PLoS Med* 2008;5(9):e193.
- [28] Xue F, Michels KB. Intrauterine factors and risk of breast cancer: a systematic review and meta-analysis of current evidence. *Lancet Oncol* 2007;8(12):1088–100.
- [29] Hoover RN, Hyer M, Pfeiffer RM, Adam E, Bond B, Chevillat AL, et al. Adverse health outcomes in women exposed in utero to diethylstilbestrol. *N Engl J Med* 2011;365:1304–14.
- [30] Innes KE, Byers TE. Preeclampsia and breast cancer risk. *Epidemiology* 1999;10(6):722–32.
- [31] Henderson TO, Amsterdam A, Bhatia S, Hudson MM, Meadows AT, Neglia JP, et al. Systematic review: surveillance for breast cancer in women treated with chest radiation for childhood, adolescent, or young adult cancer. *Ann Intern Med* 2010;152(7):444–55. W144–454.
- [32] Land CE. Studies of cancer and radiation dose among atomic bomb survivors. The example of breast cancer. *JAMA* 1995;274(5):402–7.
- [33] Nechuta S, Paneth N, Velie EM. Pregnancy characteristics and maternal breast cancer risk: a review of the epidemiologic literature. *Cancer Causes Control* 2010;21(7):967–89.
- [34] Troisi R, Doody DR, Mueller BA. A linked-registry study of gestational factors and subsequent breast cancer risk in the mother. *Cancer Epidemiol Biomarkers Prev* 2013;22(5):835–47.
- [35] Cnattingius S, Torráng A, Ekblom A, Granath F, Petersson G, Lambe M. Pregnancy characteristics and maternal risk of breast cancer. *JAMA* 2005;294(19):2474–80.
- [36] Brisken C. Progesterone signaling in breast cancer: a neglected hormone coming into the limelight. *Nat Rev Cancer* 2013;13(6):385–96.
- [37] Obr AE, Edwards DP. The biology of progesterone receptor in the normal mammary gland and in breast cancer. *Mol Cell Endocrinol* 2012;357(1–2):4–17.
- [38] Kleinberg DL, Wood TL, Furth PA, Lee AV. Growth hormone and insulin-like growth factor-I in the transition from normal mammary development to preneoplastic mammary lesions. *Endocr Rev* 2009;30(1):51–74.
- [39] Mann PC, Boorman GA, Lollini LO, McMartin DN, Goodman DG. Proliferative lesions of the mammary gland in rats, IS-2. In: *Guides for Toxicologic Pathology*. Washington, DC: STP/ARF/AFIP; 1996.
- [40] Kawaguchi H, Umekita Y, Souda M, Gejima K, Kawashima H, Yoshikawa T, et al. Effects of neonatally administered high-dose diethylstilbestrol on the induction of mammary tumors induced by 7,12-dimethylbenz[a]anthracene in female rats. *Vet Pathol* 2009;46(1):142–50.
- [41] Rothschild TC, Boylan ES, Calhoon RE, Vonderhaar BK. Transplacental effects of diethylstilbestrol on mammary development and tumorigenesis in female ACI rats. *Cancer Res* 1987;47(16):4508–16.
- [42] Muto T, Wakui S, Imano N, Nakaaki K, Takahashi H, Hano H, et al. Mammary gland differentiation in female rats after prenatal exposure to 3,3',4,4',5-pentachlorobiphenyl. *Toxicology* 2002;177(2–3):197–205.
- [43] Macon MB, Fenton SE. Endocrine disruptors and the breast: early life effects and later life disease. *J Mammary Gland Biol Neoplasia* 2013;18(1):43–61.
- [44] American Academy of Pediatrics. Breastfeeding and the use of human milk. *Pediatrics* 2012;129(3):e827–41.
- [45] Lew BJ, Collins LL, O'Reilly MA, Lawrence BP. Activation of the aryl hydrocarbon receptor during different critical windows in pregnancy alters mammary epithelial cell proliferation and differentiation. *Toxicol Sci* 2009;111(1):151–62.
- [46] Kass L, Altamirano GA, Bosquiazzo VL, Luque EH, Munoz-de-Toro M. Perinatal exposure to xenoestrogens impairs mammary gland differentiation and modifies milk composition in Wistar rats. *Reprod Toxicol* 2012;33(3):390–400.
- [47] Nebesio TD, Eugster EA. Current concepts in normal and abnormal puberty. *Curr Probl Pediatr Adolesc Health Care* 2007;37(2):50–72.
- [48] Basrur PK. Disrupted sex differentiation and feminization of man and domestic animals. *Environ Res* 2006;100(1):18–38.
- [49] Haynes BA, Mookadam F. Male gynecomastia. *Mayo Clin Proc* 2009;84:672.
- [50] Brasky TM, Li Y, Jaworowicz DJ, Potischman N, Ambrosone CB, Hutson AD, et al. Pregnancy-related characteristics and breast cancer risk. *Cancer Causes Control* 2013;24:675–1685.

- [51] Hilakivi-Clarke L, de Assis S, Warri A. Exposures to synthetic estrogens at different times during the life, and their effect on breast cancer risk. *J Mammary Gland Biol Neoplasia* 2013;18(1):25–42.
- [52] Soto AM, Brisken C, Schaeberle C, Sonnenschein C. Does cancer start in the womb? Altered mammary gland development and predisposition to breast cancer due to in utero exposure to endocrine disruptors. *J Mammary Gland Biol Neoplasia* 2013;18(2):199–208.
- [53] Hubinette A, Lichtenstein P, Ekblom A, Cnattingius S. Birth characteristics and breast cancer risk: a study among like-sexed twins. *Int J Cancer* 2001;91:248–51.
- [54] Kaijser M, Granath F, Jakobsen G, Cnattingius S, Ekblom A. Maternal pregnancy estriol levels in relation to anamnestic and fetal anthropometric data. *Epidemiology* 2000;11:315–9.
- [55] Troisi R, Potischman N, Hoover RN. Exploring the underlying hormonal mechanisms of prenatal risk factors for breast cancer: a review and commentary. *Cancer Epidemiol Biomarkers Prev* 2007;16(9):1700–12.
- [56] Titus-Ernstoff L, Hatch EE, Hoover RN, Palmer J, Greenberg ER, Ricker W, et al. Long-term cancer risk in women given diethylstilbestrol (DES) during pregnancy. *Br J Cancer* 2001;84(1):126–33.
- [57] American Cancer Society. *Breast Cancer Facts & Figures 2013–2014*. Atlanta, GA: American Cancer Society, Inc.; 2013.
- [58] Endogenous Hormones and Breast Cancer Collaborative Group. Circulating sex hormones and breast cancer risk factors in postmenopausal women: reanalysis of 13 studies. *Br J Cancer* 2011;105:709–22.
- [59] Endogenous Hormones and Breast Cancer Collaborative Group. Sex hormones and risk of breast cancer in premenopausal women: a collaborative reanalysis of individual participant data from seven prospective studies. *Lancet Oncol* 2013;14:1009–19.
- [60] Cleary MP, Grossman ME. Minireview: obesity and breast cancer: the estrogen connection. *Endocrinology* 2009;150:2537–42.
- [61] Bernstein L, Ross RK. Endogenous hormones and breast cancer risk. *Epidemiol Rev* 1993;15:48–65.
- [62] Collaborative Group on Hormonal Factors in Breast Cancer. Menarche, menopause, and breast cancer risk: individual participant meta-analysis, including 118 964 women with breast cancer from 117 epidemiological studies. *Lancet Oncol* 2012;13(11):1141–51.
- [63] Reis-Filho JS, Pusztai I. Gene expression profiling in breast cancer: classification, prognostication, and prediction. *Lancet* 2011;378(9805):1812–23.
- [64] Perou CM, Surlie T, Eisen MB, van de Rijn M, Jeffery SS, Rees CA, et al. Molecular portraits of human breast tumours. *Nature* 2000;406:747–52.
- [65] Trasande L, Attinam TM, Blustein J. Association between urinary bisphenol A concentration and obesity prevalence in children and adolescents. *JAMA* 2012;308(11):1113–21.
- [66] Sprague BL, Trentham-Dietz A, Hedman CJ, Wang J, Hemming JDC, Hampton JM, et al. Circulating serum xenoestrogens and mammographic breast density. *Breast Cancer Res* 2013;15:R45.
- [67] Chalubinski M, Kowalski ML. Endocrine disruptors—potential modulators of the immune system and allergic response. *Allergy* 2006;61(11):1326–35.
- [68] Dewitt JC, Copeland CB, Strynar MJ, Luebke RW. Perfluorooctanoic acid-induced immunomodulation in adult C57BL/6j or C57BL/6N female mice. *Environ Health Perspect* 2008;116(5):644–50.
- [69] Lopez-Espinosa M, Fletcher T, Armstrong B, Genser B, Dhataria K, Mondal D, et al. Association of perfluorooctanoic acid (PFOA) and perfluorooctane sulfonate (PFOS) with age of puberty among children living near a chemical plant. *Environ Sci Technol* 2011;45(19):8160–6.
- [70] Knox SS, Jackson T, Javins B, Frisbee SJ, Shankar A, Ducatman AM. Implications of early menopause in women exposed to perfluorocarbons. *J Clin Endocrinol Metab* 2011;96(6):1747–53.
- [71] Kavlock RJ, Daston GP, DeRosa C, Fenner-Crisp P, Gray LE, Kaattari S, et al. Research needs for the risk assessment of health and environmental effects of endocrine disruptors: a report of the U.S. EPA-sponsored Workshop. *Environ Health Perspect* 1996;104(Suppl. 4):1–26.
- [72] Zoeller RT, Brown TR, Doan LL, Gore AC, Skakkebaek NE, Soto AM, et al. Endocrine-disrupting chemicals and public health protection: a statement of principles from the endocrine society. *Endocrinology* 2012;153(9):4097–110.
- [73] Fenton SE, Reed C, Newbold RR. Perinatal environmental exposures affect mammary development, function, and cancer risk in adulthood. *Annu Rev Pharmacol Toxicol* 2012;52:455–79.
- [74] Lopez J, Ogren L, Verjan R, Talamantes F. Effects of perinatal exposure to a synthetic estrogen and progesterin on mammary tumorigenesis in mice. *Teratology* 1988;38(2):129–34.
- [75] Mori T, Bern HA, Mills KT, Young PN. Long-term effects of neonatal steroid exposure on mammary gland development and tumorigenesis in mice. *J Natl Cancer Inst* 1976;57(5):1057–62.
- [76] Mori T, Nagasawa H, Bern HA. Long-term effects of perinatal exposure to hormones on normal and neoplastic mammary growth in rodents: a review. *J Environ Pathol Toxicol* 1979;3(1–2):191–205.
- [77] Hilakivi-Clarke L, Cabanes A, de Assis S, Wang M, Khan G, Shoemaker WJ, et al. In utero alcohol exposure increases mammary tumorigenesis in rats. *Br J Cancer* 2004;90(11):2225–31.
- [78] Brown NM, Manzolillo PA, Zhang JX, Wang J, Lamartiniere CA. Prenatal TCDD and predisposition to mammary cancer in the rat. *Carcinogenesis* 1998;19(9):1623–9.
- [79] Desaulniers D, Leingartner K, Russo J, Perkins G, Chittim BG, Archer MC, et al. Modulatory effects of neonatal exposure to TCDD, or a mixture of PCBs, p,p'-DDT, and p,p'-DDE, on methylnitrosourea-induced mammary tumor development in the rat. *Environ Health Perspect* 2001;109(7):739–47.
- [80] Jenkins S, Rowell C, Wang J, Lamartiniere CA. Prenatal TCDD exposure predisposes for mammary cancer in rats. *Reprod Toxicol* 2007;23(3):391–6.
- [81] Jenkins S, Raghuraman N, Eltoum I, Carpenter M, Russo J, Lamartiniere CA. Oral exposure to bisphenol A increases dimethylbenzanthracene-induced mammary cancer in rats. *Environ Health Perspect* 2009;117:910–5.
- [82] Hilakivi-Clarke L, Cho E, Clarke R. Maternal genistein exposure mimics the effects of estrogen on mammary gland development in female mouse offspring. *Oncol Rep* 1998;5(3):609–16.
- [83] Hilakivi-Clarke L, Cho E, Onojafe I, Raygada M, Clarke R. Maternal exposure to genistein during pregnancy increases carcinogen-induced mammary tumorigenesis in female rat offspring. *Oncol Rep* 1999;6(5):1089–95.
- [84] IARC. *Pharmaceuticals. Diethylstilbestrol*. A review of human carcinogens. *IARC Monogr Eval Carcinog Risks Hum* 2012;100A:175–218.
- [85] Acevedo N, Davis B, Schaeberle CM, Sonnenschein C, Soto AM. Perinatally administered bisphenol A acts as a mammary gland carcinogen in rats. *Environ Health Perspect* 2013;121(9):1040–6.
- [86] Kociba RJ, Keyes DG, Beyer JE, Carreon RM, Wade CE, Dittenber DA, et al. Results of a two-year chronic toxicity and oncogenicity study of 2,3,7,8-tetrachlorodibenzo-p-dioxin in rats. *Toxicol Appl Pharmacol* 1978;46(2):279–303.
- [87] Wang T, Gavin HM, Arlt VM, Lawrence BP, Fenton SE, Medina D, Vorderstrasse BA. Aryl hydrocarbon receptor (AhR) activation during pregnancy, and in adult nulliparous mice, delays the subsequent development of DMBA-induced mammary tumors. *Int J Cancer* 2011;128(7):1509–23.
- [88] Warri A, Saarinen NM, Makela S, Hilakivi-Clarke L. The role of early life genistein exposures in modifying breast cancer risk. *Br J Cancer* 2008;98(9):1485–93.
- [89] Cohn BA, Wolff MS, Cirillo PM, Sholtz RI. DDT and breast cancer in young women: new data on the significance of age at exposure. *Environ Health Perspect* 2007;115(10):1406–14.
- [90] Euling SY, Selevan SG, Pescovitz OH, Skakkebaek NE. Role of environmental factors in the timing of puberty. *Pediatrics* 2008;121(Suppl. 3):S167–71.
- [91] Ojeda SR, Roth C, Mungenast A, Heger S, Mastronardi C, Parent AS, et al. Neuroendocrine mechanisms controlling female puberty: new approaches, new concepts. *Int J Androl* 2006;29(1):256–63, discussion 286–90.
- [92] Golub MS, Collman GW, Foster PM, Kimmel CA, Rajpert-De Meyts E, Reiter EO, et al. Public health implications of altered puberty timing. *Pediatrics* 2008;121(Suppl. 3):S218–30.
- [93] Vandenberg LN, Colborn T, Hayes TB, Heindel JJ, Jacobs DR, Lee DH, et al. Hormones and endocrine-disrupting chemicals: low-dose effects and non-monotonic dose responses. *Endocr Rev* 2012;33(3):378–455.
- [94] Enoch RR, Stanko JP, Greiner SN, Youngblood GL, Rayner JL, Fenton SE. Mammary gland development as a sensitive end point after acute prenatal exposure to an atrazine metabolite mixture in female Long-Evans rats. *Environ Health Perspect* 2007;115:541–7.
- [95] White SS, Stanko JP, Kato K, Calafat AM, Hines EP, Fenton SE. Gestational and chronic low-dose PFOA exposures and mammary gland growth and differentiation in three generations of CD-1 mice. *Environ Health Perspect* 2011;119(8):1070–6.
- [96] Wetherill YB, Akingbemi BT, Kanno J, McLachlan JA, Nadal A, Sonnenschein C, et al. In vitro molecular mechanisms of bisphenol A action. *Reprod Toxicol* 2007;24:178–98.
- [97] Vandenberg LN, Wadia PR, Schaeberle CM, Rubin BS, Sonnenschein C, Soto AM. The mammary gland response to estradiol: monotonic at the cellular level, nonmonotonic at the tissue-level of organization? *J Steroid Biochem Mol Biol* 2006;101:263–74.
- [98] Wadia PR, Vandenberg LN, Schaeberle CM, Rubin BS, Sonnenschein C, Soto AM. Perinatal bisphenol A exposure increases estrogen sensitivity of the mammary gland in diverse mouse strains. *Environ Health Perspect* 2007;115(4):592–8.
- [99] Kohlerova E, Skarda J. Mouse bioassay to assess oestrogenic and anti-oestrogenic compounds: hydroxytamoxifen, diethylstilbestrol and genistein. *J Vet Med A: Physiol Pathol Clin Med* 2004;51(5):209–17.
- [100] Rayner JL, Enoch RR, Fenton SE. Adverse effects of prenatal exposure to atrazine during a critical period of mammary gland growth. *Toxicol Sci* 2005;87(1):255–66.
- [101] Rayner JL, Wood C, Fenton SE. Exposure parameters necessary for delayed puberty and mammary gland development in Long-Evans rats exposed in utero to atrazine. *Toxicol Appl Pharmacol* 2004;195:23–34.
- [102] White SS, Calafat AM, Kuklenyik Z, Villanueva L, Zehr RD, Helfant L, et al. Gestational PFOA exposure of mice is associated with altered mammary gland development in dams and female offspring. *Toxicol Sci* 2007;96(1):133–44.
- [103] Vorderstrasse BA, Fenton SE, Bohn AA, Cundiff JA, Lawrence BP. A novel effect of dioxin: exposure during pregnancy severely impairs mammary gland differentiation. *Toxicol Sci* 2004;78(2):248–57.
- [104] Matsumoto C, Miyaura C, Ito A. Dietary bisphenol A suppresses the growth of newborn pups through insufficient supply of maternal milk in mice. *J Health Sci* 2004;50(3):315–8.
- [105] Vandenberg LN, Maffini MV, Schaeberle CM, Ucci AA, Sonnenschein C, Rubin BS, et al. Perinatal exposure to the xenoestrogen bisphenol-A induces

- mammary intraductal hyperplasias in adult CD-1 mice. *Reprod Toxicol* 2008;26(3–4):210–9.
- [106] Vandenberg LN, Maffini MV, Wadia PR, Sonnenschein C, Rubin BS, Soto AM. Exposure to environmentally relevant doses of the xenoestrogen bisphenol-A alters development of the fetal mouse mammary gland. *Endocrinology* 2007;148(1):116–27.
- [107] Markey CM, Luque EH, Munoz De Toro M, Sonnenschein C, Soto AM. In utero exposure to bisphenol A alters the development and tissue organization of the mouse mammary gland. *Biol Reprod* 2001;65(4):1215–23.
- [108] NTP. Multigenerational Reproductive Toxicology Study of Genistein (CAS NO. 446-72-0) in Sprague-Dawley Rats (Feed Study). Research Triangle Park, NC: National Toxicology Program; 2008.
- [109] Poulet FM, Veneziale R, Vancutsem PM, Losco P, Treinen K, Morrissey RE. Ziracin-induced congenital urogenital malformations in female rats. *Toxicol Pathol* 2005;33(3):320–8.
- [110] Felner EI, White PC. Prepubertal gynecomastia: indirect exposure to estrogen cream. *Pediatrics* 2000;105(4):E55.
- [111] Henley DV, Lipson N, Korach KS, Bloch CA. Prepubertal gynecomastia linked to lavender and tea tree oils. *N Engl J Med* 2007;356(5):479–85.
- [112] Braunstein GD. Gynecomastia. *N Engl J Med* 1993;328(7):490–5.
- [113] Latendresse JR, Bucci TJ, Olson G, Mellick P, Weis CC, Thorn B, et al. Genistein and ethinyl estradiol dietary exposure in multigenerational and chronic studies induce similar proliferative lesions in mammary gland of male Sprague-Dawley rats. *Reprod Toxicol* 2009;28(3):342–53.
- [114] Li T, Sun L, Miller N, Nicklee T, Woo J, Hulse-Smith L, et al. The association of measured breast tissue characteristics with mammographic density and other risk factors for breast cancer. *Cancer Epidemiol Biomarkers Prev* 2005;14:343–9.
- [115] Hay ED. Extracellular matrix alters epithelial differentiation. *Curr Opin Cell Biol* 1993;5:1029–35.
- [116] Lin CQ, Bissell MJ. Multi-faceted regulation of cell differentiation by extracellular matrix. *FASEB J* 1993;7(9):737–43.
- [117] Lochter A, Bissell MJ. Involvement of extracellular matrix constituents in breast cancer. *Semin Cancer Biol* 1995;6:165–73.
- [118] Maffini MV, Soto AM, Calabro JM, Ucci AA, Sonnenschein C. The stroma as a crucial target in rat mammary gland carcinogenesis. *J Cell Sci* 2004;117(Pt 8):1495–502.
- [119] Medina D. Chemical carcinogenesis of rat and mouse mammary glands. *Breast Dis* 2007;28:63–8.
- [120] IOM. *Breast Cancer and the Environment: A Life Course Approach*. Institute of Medicine. Washington, DC: The National Academies Press; 2011.
- [121] Rasmussen SB, Young LJT, Smith GH. Preparing mammary gland whole mounts from mice. In: Ip MM, Asch BB, editors. *Methods in Mammary Gland Biology and Breast Cancer Research*. New York, NY: Kluwer Academic/Plenum Publishers; 2000. p. 75–85.
- [122] Kavanaugh-Lynch M, Francis E, Darney S, Birnbaum L, Reinlib L, Thompson C, et al. Report to funders: Mammary Gland Evaluation and Risk Assessment Workshop. In: Fenton S, Rudel R, editors. *Proceedings of the Mammary Gland Evaluation and Risk Assessment Workshop Round Robin Meeting*. 2009.
- [123] Boberg J, Mandrup KR, Jacobsen PR, Isling LK, Hadrup N, Berthelsen L, et al. Endocrine disrupting effects in rats perinatally exposed to a dietary relevant mixture of phytoestrogens. *Reprod Toxicol* 2013;40C:41–51.
- [124] Wang XJ, Bartolucci-Page E, Fenton SE, You L. Altered mammary gland development in male rats exposed to genistein and methoxychlor. *Toxicol Sci* 2006;91(1):93–103.
- [125] You L, Sar M, Bartolucci EJ, McIntyre BS, Sriperumbudur R. Modulation of mammary gland development in prepubertal male rats exposed to genistein and methoxychlor. *Toxicol Sci* 2002;66(2):216–25.
- [126] Lucas JN, Rudmann DG, Credille KM, Irizarry AR, Peter A, Snyder PW. The rat mammary gland: morphologic changes as an indicator of systemic hormonal perturbations induced by xenobiotics. *Toxicol Pathol* 2007;35(2):199–207.